

ASPEN Lipid Injectable Emulsion Safety Recommendations for Adult Patients¹

- Lipid injectable emulsions (ILEs) are intended as a daily source of energy and fatty acids (FAs) for parenteral nutrition (PN) therapy.
- There are currently four FDA approved ILE products with an adult indication available in the US. Alternative ILE products for adults are those with decreased amounts of soybean oil.

Dosing and Prescribing

- The dose of ILE should be prescribed as amount **per day** (ie, g/d or kcal/d for adults). **Table 1** provides adult daily dosing recommendations.

ILE	SO-ILE	SO,MCT,OO,FO-ILE	OO,SO-ILE
Oil Source	100% soybean oil	30% soybean oil, 30% MCT oil, 25% olive oil, 15% fish oil	80% olive oil, 20% soybean oil
Recommended Dose	Critically ill: <1g/kg/d Stable: 1 g/kg/d	1–2 g/kg/d	1–1.5 g/kg/d
Maximum Dose	2.5 g/kg/d	2.5 g/kg/d	2.5 g/kg/d
Maximum Infusion Rate	0.11 g/kg/h	0.11 g/kg/h	0.11 g/kg/h

Abbreviations: ILE, lipid injectable emulsion; OO,SO-ILE, olive oil, soybean oil ILE—Clinolipid® (Baxter Healthcare Corporation); SO,MCT,OO,FO-ILE, soybean oil, medium-chain triglycerides, olive oil, fish oil ILE—SMOFLipid® (Fresenius Kabi); SO-ILE, soybean oil ILE—Intralipid® (Baxter Healthcare Corporation), Nutrilipid® (B. Braun Medical, Inc).

- The standardized PN order should include the **brand name** of the ILE product due to the variable composition of ILE products.
- With reduced soybean oil products, ILE may be provided to critically ill adults upon PN initiation without producing the inflammatory and immunosuppressive effects associated with SO-ILE.
- Review ILE Prescribing Information for contraindications and prescribe appropriately to avoid allergic reactions. Allergy contraindications for ILEs will vary based on oil source(s) and other active ingredients and excipients.
- Adult minimum EFA recommendations are at least 2%–4% of energy from LA and 0.25%–0.5% of energy from ALA.
 - » Dosing for maintenance or long-term therapy should be about 1 g/kg/day in the stable adult patient.

- » ILEs that are less than 100% soybean oil require a higher minimum dose vs SO-ILE due to lower LA and ALA content.

Daily total caloric dose	Daily dose SO ILE (20%)		Daily dose SO,MCT, OO,FO ILE (20%)		Daily dose OO, SO ILE (20%)	
	mL*	g	mL*	g	mL*	g
1000	20.4	4.1	62.9	12.6	61.5	12.3
1500	30.6	6.1	94.3	18.9	92.2	18.4
2000	40.8	8.2	125.7	25.1	122.9	24.6
Daily dose, % total kcal to meet ≥ 2% of total kcal from LA	≥ 7%		≥ 12.5%		≥ 12.5%	

Abbreviations: ALA, α-linolenic acid; EFA, essential fatty acid; ILE, lipid injectable emulsion; LA, linoleic acid; OO,SO-ILE, olive oil, soybean oil ILE—Clinolipid® (Baxter Healthcare Corporation); SO,MCT,OO,FO-ILE, soybean oil, medium-chain triglycerides, olive oil, fish oil ILE—SMOFLipid® (Fresenius Kabi); SO-ILE, soybean oil ILE—Intralipid® (Baxter Healthcare Corporation), Nutrilipid® (B. Braun Medical, Inc).

ILE LA and ALA Content: ²⁻⁵ SO ILE contains 88-124 mg/mL LA and 8-22 mg/mL ALA; SO,MCT,OO,FO ILE contains 35 mg/mL LA (mean) and 4.5 mg/mL ALA (mean); OO,SO ILE contains 35.8 mg/mL LA (mean) and 4.7 mg/mL ALA (mean)

Calculations are based on the ILE content (mean or range) for LA as an amount required to provide at least 2% of total kcal from LA as reported for the products * When dispensing ILE, round up to the nearest mL

- The minimum dose to prevent EFAD is **NOT** the daily recommended dose.
- EFA content may vary within a preestablished range which may account for variations of reported amounts in the literature.
- These are minimum estimated dosing recommendations; individuals may require higher ILE dosing to prevent or treat EFAD depending on clinical circumstances.
- Greater than ILE minimum dosing allows for a reduction in the amount of dextrose needed to provide appropriate energy intake.

Continued ►


Monitoring

- Check serum triglycerides at baseline and weekly for hospitalized patients and monthly for long-term PN patients.
- Hold or limit ILE if triglyceride level is > 400 mg/dL. Note, withholding ILE greater than 2 weeks is not advised due to risk for EFAD.

Stability/Compatibility

Total Nutrient Admixture (TNA) Stability

- The stability and compatibility of TNA is dependent on many factors including, electrolyte content and specific salts used, final concentration of macronutrients, and pH (These are not the same across all manufacturer's products).
- A pharmaceutical review should be conducted by a knowledgeable pharmacist to assess the stability and compatibility of ILE with the other components of the PN admixture.
- ILE manufacturers should be contacted to determine the acceptable range of macro- and micronutrient concentrations for TNA stability.
- **Figure 1** illustrates phases of TNA physical destabilization:



Adapted with permission from Pertkiewicz M et al.⁸ Copyright © European Society for Clinical Nutrition and Metabolism. Published by Elsevier B.V. All rights reserved.

These recommendations do not constitute medical or other professional advice and should not be taken as such. To the extent that the information published herein may be used to assist in the care of patients, this is the result of the sole professional judgment of the attending healthcare professional whose judgment is the primary component of quality medical care. The information presented is not a substitute for the judgment by the healthcare professional. Circumstances in clinical settings and patient indications may require actions different from those recommended in this document, and in those cases, the judgment of the treating professional should prevail. ASPEN does not endorse any particular brand of products mentioned herein.


References

- ¹ Mirtalio JM, Ayers P, Boullata J, et al. ASPEN Lipid Injectable Emulsion Safety Recommendations, Part 1: Background and Adult Considerations. *Nutr Clin Pract.* 2020;35(5):769-782.
- ² Intralipid 20% (IV fat emulsion) Prescribing Information. In: Fresenius Kabi; 2015.
- ³ Nutrilipid 20% (lipid injectable emulsion) Prescribing Information. In: B. Braun Medical Inc.; 2014.

Preparation & Administration

- When ILE is ordered as part of a TNA, it may be infused over 24 hours. If ordered as a separate infusion, it should be infused over a maximum of 12 hours. Can split into two 12 hour doses if needed for volume limitations.
- ILE should be administered using DEHP-free tubing due to concern of DEHP plasticizer leaching into ILE.
- A 1.2 micron filter should be used for a TNA formulation or as a separate ILE infusion. See **Figure 2** for appropriate set-up.⁹
- Administration of medications through the PN venous catheter should be avoided due to increased risk for catheter related blood stream infection (CRBSI) and compatibility concerns.
 - » ILE manufacturers should be contacted for medication compatibility information.

FIGURE 2. APPROPRIATE SET-UP OF PN WITH FILTER



Reprinted from Worthington, et al.⁹ Copyright © ASPEN

- ⁴ Clinolipid 20% (lipid injectable emulsion) Prescribing Information. In: Baxter Healthcare Corporation; 2016.
- ⁵ SMOFlipid (lipid injectable emulsion) Prescribing Information. In: Fresenius Kabi LLC USA; 2020.
- ⁶ McClave SA, Taylor BE, Martindale RG, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr.* 2016;40(2):159-211.
- ⁷ Ayers P, Bobo E, Hurt RT, Mays A, Worthington P. *ASPEN Parenteral Nutrition Handbook, 3rd Edition.* 3rd ed. Silver Spring, Maryland: ASPEN; 2020.
- ⁸ Pertkiewicz M, Cosslett A, Mühlebach S, Dudrick SJ. Basics in clinical nutrition: Stability of parenteral nutrition admixtures. *e-SPEN, the European e-Journal of Clinical Nutrition and Metabolism.* 2009;4(3):e117-e119.
- ⁹ Worthington P, Gura KM, Kraft MD, et al. Update on the Use of Filters for Parenteral Nutrition: An ASPEN Position Paper. *Nutr Clin Pract.* 2021;36(1):29-39.

This educational sheet has been supported by:

